

EARTH: 15 MILLION YEARS AGO

Masataka Mizushima

Department of Physics, University of Colorado
Boulder, Colorado 80309

Abstract

In Einstein's general relativity theory the metric component g_{xx} in the direction of motion (x-direction) of the sun deviates from unity due to a tensor potential caused by the black hole existing around the center of the galaxy. Because the solar system is orbiting around the galactic center at 200 km/s, the theory shows that the Newtonian gravitational potential due to the sun is not quite radial. At the present time, the ecliptic plane is almost perpendicular to the galactic plane, consistent with this modification of the Newtonian gravitational force. The ecliptic plane is assumed to maintain this orientation in the galactic space as it orbits around the galactic center, but the rotational angular momentum of the earth around its own axis can be assumed to be conserved. The earth is between the sun and the galactic center at the summer solstice all the time. As a consequence, the rotational axis of the earth would be parallel to the axis of the orbital rotation of the earth 15 million years ago, if the solar system has been orbiting around the galactic center at 200 km/s. The present theory concludes that the earth did not have seasons 15 million years ago. Therefore, the water on the earth was accumulated near the poles as ice and the sea level was very low. Geological evidence exist that confirms this effect. The resulting global ice-melting started 15 million years ago and is ending now.

Subect heading: General Relativity, Solar System, Milky Way Galaxy

If we take

$$g_{ij} = g_{ij}^{(\circ)} + h_{ij}, \quad (1)$$

where

$$g_{00}^{(\circ)} = 1, \quad g_{\alpha 0}^{(\circ)} = 0, \quad g_{\alpha\beta}^{(\circ)} = -\delta_{\alpha\beta}, \quad (2)$$

$\alpha = 1, 2, 3$, for the general relativistic metric

$$(ds)^2 = g_{ij} dx^i dx^j, \quad (3)$$

and assume that the h_{ij} are all so small that their powers higher than the first are negligible, Einstein's general relativistic field equations can be simplified into those of the linearized Einstein theory.

Introducing

$$h_{ij} = \phi_{ij} - \frac{1}{2} \phi g_{ij}^{(\circ)} \quad (4)$$

where $\phi = \phi_i^i$, and $\phi_j^i = g^{(\circ)ik} \phi_{kj}$, and assuming that

$$\frac{\partial \phi_j^i}{\partial x^i} = 0, \quad (5)$$

we see that Einstein's field equation can be approximated as

$$\left(\frac{\partial^2}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \phi_j^i = \frac{16\pi G}{c^4} T_j^i. \quad (6)$$

In eq. (6), T_j^i is a component of the mass energy-momentum tensor such that $T_0^0 = \mu c^2$.

It is easy to solve eq. (6), because it is mathematically identical to the Maxwell field equations, to obtain

$$\phi_{00} = -\Sigma_a \frac{4GM_a}{c^2 r_a} \big|_{t'}, \quad (7a)$$

$$\phi_{\alpha 0} = \Sigma_a \frac{4GM_a v_{a,\alpha}}{c^3 r_a} \big|_{t'}, \quad (7b)$$

$$\phi_{\alpha\beta} = -\Sigma_a \frac{4GM_a v_{a,\alpha} v_{a,\beta}}{c^4 r_a} \big|_{t'}, \quad (7c)$$

where M_a is the mass of a point source located at a distance r_a from the observation point, and $v_{a,\alpha}$ is the α -component of its velocity. Because eq. (6) is linear in μ , the mass density, we see that the right-hand sides of eqs. (7a) through (7c) are given by the sum over existing point masses M_a . Quantities on the right-hand side of eqs. (7a) through (7c) are to be evaluated at the retardation time t' , but the retardation effect can be

neglected.

In the following discussion, we take the solar mass as M_s and the black hole mass as M_b , which exists near the center of the Milky way galaxy. In the Newtonian limit, $v/c \rightarrow 0$, we see that

$$g_{00} = -1 + \frac{1}{2}\phi_{00} = -1 - \frac{2GM_s}{c^2 r_s} - \frac{2GM_b}{c^2 r_b} \quad (8)$$

where r_s and r_b are the distances to the sun and to the galactic center, respectively. The terms proportional to M_s and M_b give the Newtonian potentials due to the sun and the black hole at the galactic center, respectively. The Newtonian potential due to M_b is negligible compared to that due to the sun. In the solar system, the solar mass M_s is not moving, but the source mass M_b is moving in the direction tangential to the galactic center, which we call the x -direction, so that $\phi_{yy} = \phi_{zz} = 0$, and

$$g_{xx} - g_{yy} = g_{xx} - g_{zz} = \frac{1}{2}\phi_{xx} = -\frac{2GM_b v_g^2}{c^4 r_b} \equiv -P, \quad (9)$$

where v_g is the velocity of the galactic center with respect to the sun, in the direction tangential to the galactic center. Actually, we can write this anisotropy P as

$$P = \frac{2GM_b v_g^2}{c^4 r_b} = 2(v_g/c)^4, \quad (10)$$

using the relation $GM_b/r_b = v_g^2$, for a circular orbit. Because the orbiting speed v_g is 200 km/s, the anisotropy is $P = 2 \times 10^{-12}$ at the solar system. The retardation effect is neglected in this estimation.

Einstein also proposed an equation of motion under gravity:

$$\frac{d^2 x^i}{ds^2} = -\Gamma_{jk}^i \frac{dx^j}{ds} \frac{dx^k}{ds}, \quad (11)$$

where

$$\Gamma_{jk}^i = \frac{1}{2}g^{im} \left(\frac{\partial g_{mk}}{\partial x^j} + \frac{\partial g_{mj}}{\partial x^k} - \frac{\partial g_{kj}}{\partial x^m} \right) \quad (12)$$

is a component of the Christoffel symbol. In the nonrelativistic limit, $s = ct$, and the first term is obtained by taking $j = k = 0$ in eq. (11), so that

$$\frac{d^2 x}{dt^2} \approx -\Gamma_{00}^1 = \frac{c^2}{2} g^{xx} \frac{\partial g_{00}}{\partial x}. \quad (13)$$

To the first order in P , we obtain

$$g^{xx} = 1 - \frac{1}{2}\phi_{xx} = 1 + P, \quad g^{yy} = g^{zz} = 1. \quad (14)$$




Figure 1: At present the ecliptic plane is almost perpendicular to the orbiting velocity $\mathbf{v}_g$ of the solar system around the galactic center; the ecliptic plane contains the galactic center in such a way that the earth is between the galactic center and the sun at the summer solstice, which implies that the rotational angular momentum of the earth, $\mathbf{S}$, is in the galactic plane, about $\theta = 23.4^\circ$ with respect to $\mathbf{v}_g$ at present. Assuming that the orbiting speed, 200 km/s, remained constant with the radial distance 3×10^{20} m from the galactic center, orbiting velocity $\mathbf{v}_g$ was 23.4° off from the present direction 15 million years ago. Therefore, $\mathbf{v}_g$ was parallel to $\mathbf{S}$ 15 million years ago.

neglecting ϕ_{00} compared to 1. Therefore, eq. (13) reduces to

$$\frac{d^2 \mathbf{r}}{dt^2} = -(\nabla + (P\mathbf{v}_g/v_g^2)(\mathbf{v}_g \cdot \nabla)) \frac{GM_s}{r_s}, \quad (15)$$

where M_s is the solar mass. In eq. (7a), ϕ_{00} is the sum of the Newtonian potentials due to the galactic center and to the sun, but the contribution to eq. (15) due to the galactic center is negligible. The extra term, proportional to P , in eq. (15), provide the torque to keep the ecliptic plane containing the galactic center during the orbital motion of the solar system around the galactic center (Mizushima., 1992)

Because the solar system is orbiting around the galactic center, the Newtonian gravitational force due to the sun is not isotropic, but anisotropic in the direction of the orbiting velocity around the galactic center. The present orientation of the ecliptic plane is almost in the direction that contains the galactic center, as shown in Fig. 1, and that can be understood taking this general relativistic anisotropy of the Newtonian gravitational force into account.

According to the present theory, the anisotropy of the space force kept the orientation of the ecliptic plane perpendicular to the $\mathbf{v}_g$ vector. Assuming that the orbiting speed has been 200 km/s, the vector $\mathbf{r}_b$ 15 million years ago was off by 23.4° from the present direction of $\mathbf{r}_b$. Another interesting situation of the present orbit of the earth is that the sun is between Earth and the galactic center at the winter solstice, and the earth is between the sun and the galactic center at the summer solstice. Thus, the rotational angular momentum of the earth is in the galactic plane making an angle of 23.4° away from the galactic center. Therefore, 15 million years ago, the orbital angular momentum of the earth around the sun was parallel to its rotational angular momentum $\mathbf{S}$ (Fig.1).

At present, the sun is at 23.4° North at the summer solstice and 23.4° South at the winter solstice. Analyzing the reported average temperature at a given location on Earth, we see that the freezing temperature at that location appears when the angular distance of the ecliptic plane from the vertical of that location is more than 50° at sea level and 40° in the high mountains. The average temperature at each location is mostly dependent on the rate of energy input by the solar radiation, which is proportional to $\cos^2 \theta$ where θ is the angular distance between the sun at noon and a vertical at the observational location on the earth. (Mizushima, 2008a).

If we assume that solar radiation was the main source of the variable energy input to the surface of the earth 15 million years ago, as it is now, the above observation means that the earth everywhere with a latitude greater than 40° , both North and South, stayed at a temperature below freezing year around 15 million years ago because there were no seasons. The equatorial region stayed warm, and water kept evaporating from the oceans. But atmospheric water vapor fell down to the surface of the earth as snow at higher latitude regions both North and South. On the continents, the snow did not melt, and accumulated as ice. Although ice also evaporates, it does so very slowly. Thus, a considerable fraction of earth's water remained as ice during the period around 15 million years ago, which resulted in very low sea level. The geological evidence to confirm this effect exists (Vail, et al., 1977), as shown in Fig. 2.




Figure 2: Geological evidence that sea level was at its lowest 15 million years ago (Veil, et al., 1982).

As the ecliptic plane began to deviate from the equator, temperatures at critical latitudes, both North and South, rose, and ice started to melt in the summer seasons, and will keep melting for some time to come.

It is found (Mizushima, 2008b) that the tangential velocity of the solar system around the galactic center, the orbiting speed v_g as cited in eq. (9) and Fig. 1, was very much smaller than the present value 200 km/s during the earlier life of the solar system. If that is true, the anisotropy factor P was much smaller than its present value. Therefore, the anisotropy effect shown by eq. (15) appeared only during the very recent period as discussed here.

References

- Mizushima, M., 1992, J. Phys. Soc. Japan **61**, 2673
 Mizushima, M., 2008a, Proc. Dynamic Sys. Appl. **5**, 339
 Mizushima, M. 2008b, arXiv 0801.4809
 Vail, P. R., Mitchem, R. M. Jr., and Thomson, S. III, 1977, *Seismic Stratigraphy. Application to Hydrocarbon Exploration*, Memoir 26 p.85, reproduced on p.278 of *Marine Geology*, ed. Kennett, J. P. Prentice Hall, Englewood, NJ (1982).